



ELSEVIER

Developmental Brain Research 135 (2002) 105

DEVELOPMENTAL
BRAIN
RESEARCH

www.elsevier.com/locate/brain

Author Index

- Avivi, C., Cui, S., Goren, S. and Goldstein, R.S.
Differences in neuronal differentiation between the transient cranial (Frorieps') and normal dorsal root ganglia (135) 19
- Bothwell, M., see Karam, S.D. (135) 29
- Boyd, A.W., see Karam, S.D. (135) 29
- Brown, R.E., see Penner, M.R. (135) 71
- Carrey, N., see Penner, M.R. (135) 71
- Cottriall, C.L., see Vessey, K.A. (135) 79
- Cui, S., see Avivi, C. (135) 19
- de Rogalski Landrot, I., see Riviello, P. (135) 101
- Dottori, M., see Karam, S.D. (135) 29
- Fritzsche, B., see Maklad, A. (135) 1
- Goldstein, R.S., see Avivi, C. (135) 19
- Goren, S., see Avivi, C. (135) 19
- Haber, S., see Lapidot, A. (135) 87
- Henderson, J.T., see Karam, S.D. (135) 29
- Holmes, G.L., see Riviello, P. (135) 101
- Hongo, S., see Nakada, N. (135) 45
- Hongo, S., see Ohki, T. (135) 55
- Hutchinson, I., see Temple, E.C. (135) 65
- Jinks, A.L., see Temple, E.C. (135) 65
- Karam, S.D., Dottori, M., Ogawa, K., Henderson, J.T., Boyd, A.W., Pasquale, E.B. and Bothwell, M.
EphA4 is not required for Purkinje cell compartmentation (135) 29
- Kuhn, H.-J., see Malz, C.R. (135) 39
- Laing, D.G., see Temple, E.C. (135) 65
- Lapidot, A. and Haber, S.
Effect of endogenous β -hydroxybutyrate on brain glucose metabolism in fetuses of diabetic rabbits, studied by ^{13}C magnetic resonance spectroscopy (135) 87
- Maeda, A., see Nakada, N. (135) 45
- Maeda, A., see Ohki, T. (135) 55
- Maklad, A. and Fritzsche, B.
The developmental segregation of posterior crista and saccular vestibular fibers in mice: a carbocyanine tracer study using confocal microscopy (135) 1
- Malz, C.R. and Kuhn, H.-J.
Calretinin and FMRFamide immunoreactivity in the nervous terminalis of prenatal tree shrews (*Tupaia belangeri*) (135) 39
- McBrien, N.A., see Vessey, K.A. (135) 79
- McFadyen, M.P., see Penner, M.R. (135) 71
- Nakada, N., Hongo, S., Ohki, T., Maeda, A. and Takeda, M.
Molecular characterization of NDRG4/Bdm1 protein isoforms that are differentially regulated during rat brain development (135) 45
- Nakada, N., see Ohki, T. (135) 55
- Ogawa, K., see Karam, S.D. (135) 29
- Ohki, T., Hongo, S., Nakada, N., Maeda, A. and Takeda, M.
Inhibition of neurite outgrowth by reduced level of NDRG4 protein in antisense transfected PC12 cells (135) 55
- Ohki, T., see Nakada, N. (135) 45
- Pasquale, E.B., see Karam, S.D. (135) 29
- Penner, M.R., McFadyen, M.P., Pinaud, R., Carrey, N., Robertson, H.A. and Brown, R.E.
Age-related distribution of c-fos expression in the striatum of CD-1 mice after acute methylphenidate administration (135) 71
- Pinaud, R., see Penner, M.R. (135) 71
- Riviello, P., de Rogalski Landrot, I. and Holmes, G.L.
Lack of cell loss following recurrent neonatal seizures (135) 101
- Robertson, H.A., see Penner, M.R. (135) 71
- Takeda, M., see Nakada, N. (135) 45
- Takeda, M., see Ohki, T. (135) 55
- Temple, E.C., Hutchinson, I., Laing, D.G. and Jinks, A.L.
Taste development: differential growth rates of tongue regions in humans (135) 65
- Vessey, K.A., Cottriall, C.L. and McBrien, N.A.
Muscarinic receptor protein expression in the ocular tissues of the chick during normal and myopic eye development (135) 79



